



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

DIOPHANTINE ANALYSIS.

86. Proposed by B. F. FINKEL, A. M., M.Sc., Professor of Mathematics and Physics, Drury College, Springfield, Mo.

Prove that the congruence $x^2 = 1457 \equiv 0 \pmod{2389}$ is not possible.

A solution of this problem is given in Mathews' *Theory of Numbers*, page 42. Ed.

87. Proposed by LON C. WALKER, A.M., Professor of Mathematics, Petaluma High School, Petaluma, Cal.

Find three numbers in arithmetical progression the sum of whose cubes is a cube.

Solution by the PROPOSER.

Let tx , ty , tz represent the three required numbers. Then we have

$$t^3x^3 + t^3y^3 + t^3z^3 = w^3; \text{ or, } x^3 + y^3 + z^3 = w^3/t^3 = v^3 \dots (1), \text{ assume.}$$

Transposing (1), we have to satisfy the equation

$$v^3 - x^3 - y^3 = z^3 \dots (2).$$

Put $x = a(r-q)$, $y = aq$, $z = a[p - (rq^2/p^2)]$, and $v = ap$. Then we have, after dividing by a^3 ,

$$p^3 - (r-q)^3 - q^3 = (p - \frac{rq^2}{p^2})^3 \dots (3).$$

Whence, by involution and reduction,

$$r = \frac{3p^3q}{p^3 + q^3}.$$

Now take $a = p^3 + q^3$, and we have

$$x = q(2p^3 - q^3), y = q(p^3 + q^3), z = p(p^3 - 2q^3), v = p(p^3 + q^3),$$

where p and q may have any values subject to the condition $p^3 > 2q^3$.

If we take $p=2$, $q=1$, after dividing all by 3, we have, $3^3 + 4^3 + 5^3 = 6^3$, the least three positive integral numbers that satisfy conditions of the problem.

[See *Mathematical Magazine*, page 154, from whence the above solution was copied.]

88. Proposed by L. C. WALKER, A. M., Professor of Mathematics, Petaluma High School, Petaluma, Cal.

Find three square numbers in harmonical progression.

Solution by HARRY S. VANDIVER, Bala, Pa.

Solutions of this problem have been given on pages 82-83 of Vol. VII, of the MONTHLY, substantially as follows: